

R E M A R K S

O N

781. e. 23
5

Dr. BOERHAAVE'S

T H E O R Y

O F T H E

A T T R I T I O N o f t h e B L O O D

I n t h e L U N G S

B Y

S A M U E L M U S G R A V E, M. A.

O f *Corpus-Christi-College*, OXFORD.

L O N D O N,

Printed in the Year MDCCLIX.



THE celebrated Dr. BOERHAAVE in his *Institutiones Medicæ*, lays it down as a fundamental Principle, that the Blood, in passing through the Lungs, undergoes a Degree of Attrition considerably greater than in any other Part of the Body. This Attrition, according to his System, eminently increases the Heat of the Blood, at the same Time that it unites the elementary Parts of it more intimately, and makes the whole Mass more dense and compact.

I am so far from assenting to this Theory, that I cannot but apprehend it is abundantly confuted by the Texture and Appearance of the Parts themselves. Is it to be supposed, for Instance, that the right Ventricle of the Heart, which is thinner and apparently weaker than the left, can communicate a greater Impulse to the Blood? This however is one of the first Requisites in producing a greater Degree of Attrition. There is still a more remarkable Difference between the Pulmonary Artery, and the Aorta; the one, being loose and flaccid; the other, of so dense and compact a Nature, as to resemble, in some Measure, a Cartilage.

If we go on to examine the Texture of the Parts, through which these Circulations are respectively performed, and observe how tender and spongy the Lungs are, in Comparison of the Muscular Parts of the Body, we cannot avoid coming to the same Conclusion; unless we would suppose that Nature made the stronger Substance to resist the weaker Impression: which surely would be interpreting its Meaning very perversely.

Neither is it material to say that the Blood may suffer considerable Attrition by the alternate Motion of the Lungs in Respiration; either when expanded by the Air, or compressed by the Thorax. For, Action and Reaction being equal, it makes no Difference in the Effect, whether the Lungs be every Moment pressed upon the Blood, or the Stream of Blood forced against the Lungs. Should we therefore admit the Sum of the Attritions produced by these two Causes in the Lungs, to be greater than that which subsists in the corresponding Arteries of the Body, the same Objection would still remain, in regard to the Structure of the Parts. It would still be impossible to comprehend how the Author of Nature could proportion his Work so ill, as to make the Power of Resistance greatest, where the Impression was least: to give an unnecessary Degree of Strength to one System of Arteries, or an insufficient one to the other. The

The Difficulties that attend this Theory are not more extraordinary, than the Proofs, by which it is supported, are defective. In regard to the Velocity of the two Circulations, Dr. HALLER expresses himself thus: *Cum sanguinis tantum per pulmones transeat in dato tempore, quantum per universum corpus humanum, sequitur, ut sanguis per pulmonem in eadem ratione fluat celerius, quàm ipse pulmo minor est.* (In Boerhaav. *Prælection. Acad.* Vol. II. p. 169.) It is hard to say whether the Doctor is more unfortunate here in his Premises, or his Conclusion. In the first Place, that equal Quantities of Blood pass in equal Times through the Lungs, and through the larger Circuit of the Body, is asserted, as far as I can inform myself, without any manner of Foundation either in Reason or Experiment. And secondly, admitting it to be true, the Consequence would be just the contrary of what the Doctor supposes; namely, that the Motion of the Blood through the Lungs must be slower, and in the very same Proportion too, that he assigns for its being quicker. In a late Edition of the Commentary on *Boerhaave's* Institutions, I find the following Note, by which it appears that the Doctor begins to suspect at least, the Justness and Force of his Reasoning: *Hæc omnia expedita sunt. Unum superest an velocius trajicitur per pulmones Sanguis? Videtur etiam lentius*

lentiùs trajici, quia sanguis per corpus trajectus viam multò majorem eodem tempore absolvit, — & video hanc rationem fusè & ornatè proponi a Cl. Krugero in Physiologia Germanicè scriptâ. Vol. II. p. 230. Ed. Gotting. 1745.

It has indeed been observed by Dr. Hales (*Hæmostat. p. 67.*) that in the Lungs of a Frog, the Blood moves with greater Velocity than in any other Part of the Body. We are not however to make any general Inferences from this Phænomenon, which seems to depend upon the peculiar Structure of the Frog and some other small Animals, the Heart of which, having only one Ventricle, must of Necessity give the same Impulse to all the circulating Blood. We cannot, I say, conclude from Instances of this Kind, that the relative Velocity of the two Circulations is exactly the same in larger Animals, where the Heart, having two Ventricles, may possibly impel the Blood with different Degrees of Force.

But to consider this Experiment in another Light. It appears that the Power exerted at the Origin of the pulmonary Artery and the Aorta being equal, the Blood moves with greater Velocity through the Lungs, than through any other Part of the Body: what are we to conclude from this,
but

but that the Blood-Vessels of the Lungs are larger, allowing the Blood to pass through them with greater Freedom? Which being once admitted, Dr. *Boerhaave's* Theory is directly overthrown, in the very Instance that Dr. *Haller* refers to in Support of it (Vol. II. p. 230.) and still more clearly with respect to larger Animals, where the Force impelling the Blood into the Lungs is with good Reason supposed to be less than that which throws it into the Aorta.

F I N I S.

...the Blood Vessels of the Lungs are larger
allowing the Blood to pass through them with
greater Freedom? Which being once admitted
Dr. Barrow's Theory is greatly overthrown
the very Inference that Dr. Barrow refers to is
part of it (Vol. II. p. 111) and more
with respect to the Lungs which the
impelling the Blood is with
Reason supposed to be the same as
it into the Arteries.



BRITAN